

Enhanced skin structure through innovative anti-aging treatment with Intense Pulsed Light (IPL) and boswellia nanoparticles

published in *Ästhetische Dermatologie* 2009 (4), 28-33

[IPL-activated boswellia-(frankincense-)Triterpenes]

Boswellia nanoparticles (Boswellia Carteri, Boswellia sacra)

Frankincense is a resin gained by incising the barks of desert trees of the species boswellia. Key growing areas of the boswellia trees are the Middle East specifically Oman, Yemen, Somalia and India. The escaping resin hardens in the sun and after a laborious manual harvesting with a specific scraping knife it is sold in frankincense bazaars. Frankincense extracts have anti-inflammatory properties and are rather effective against allergies and tumors. They are successful remedies against inflammatory skin diseases as for instance actinic keratoses, psoriasis and acne.

According to the current state of knowledge, the pharmacologically active ingredients of frankincense resins are the boswellic acids. Sashwati et al. [DNA and Cell Biology 24 (4), 244-255, 2005] could prove the anti-inflammatory and collagen protecting mechanism of acetyl-keto-boswellic acid which consists of inhibiting the matrix metalloproteinases (MMP).

To treat inflammatory skin diseases standardized frankincense extract containing at least 30 % acetyl-keto-boswellic acids was isolated and embedded in nanoparticles. This active agent concentrate is only useful for the skin after encapsulating it into nanoparticles since frankincense extract itself has strongly adhesive properties and cannot penetrate into the skin. Nanoparticles are spherical structures with 60 to 100 nm in size and with a shell consisting of the natural phospholipid of the skin, the so-called phosphatidylcholine (PC). Nanoparticles may encapsulate lipophilic substances which then form watery dispersions. Once applied on the skin the PC-shell of the nanoparticles fuses with the stratum corneum of the skin while the content of the nanoparticles is transported into the deeper skin layers. Without this specific transport mechanism certain active agents cannot pass the skin barrier.

The boswellia nanoparticles can be incorporated into a DMS[®] cream (Derma Membrane Structure, available at KOKO, Leichlingen). The result is a barrier strengthening skin care cream rich in active agents which is used to treat inflammatory and proliferating skin diseases.

In a 1:1 ratio (gel : active agent) boswellia nanoparticles can be mixed into a specific cosmetic gel base (KOKO, Leichlingen) which is free of preservatives. This boswellia gel was used for the treatments in the present study.

Pre-studies with boswellia nanoparticles

Boswellia nanoparticles for the treatment of inflammatory skin diseases

1 – 5 % boswellia nanoparticles combined with a topical DMS[®] cream were applied twice a day, morning and evening, together with a slight massage of the affected skin areas. After one week of applications already, actinic keratosis, neurodermatitis and psoriasis inflammations could be visibly reduced in pilot studies. After a treatment period of about 6 weeks three patients out of a group of five were completely healed of inflammatory skin diseases. The result has been histologically proved in one patient. The symptoms of 2 patients have considerably improved.

Intense pulsed light (DepiLight IPL)-System

IPL is an abbreviation for intense pulsed light and refers to a specific technology to remove unwanted hair and to modify the skin structure with a high energy light source. In contrast to laser technology which is based on a singular and constant wave length as e.g. 810 nm for laser diodes, the IPL technology uses the complete spectrum of a xenon light source which then is reduced to the required

Test person (female)	Skin condition before the treatment	Skin condition after the treatment	Subjective evaluation	Energy density / pulse	Dermatoscopy
B.N. 20.07.60	pre-aged skin with deep wrinkles	noticeably firmer skin, reduction of wrinkles	considerable improvement: +++	8 J* cm ² / 2 pulses	reduction of wrinkle depth, improved skin barrier
G.R. 12.04.39	signs of aging skin, barrier disorders	improved skin condition, firming, improved skin barrier	visible improvement, rejuvenation: +++	9 J* cm ² / 1 pulse	firming, improved skin barrier, reduced wrinkle depth
M.M. 05.09.43	aged skin, wrinkles, dry skin, barrier disorders	firming, reduction of wrinkles, improved skin hydration	Visible improvement: ++	11 J* cm ² / 1 pulse	improved skin barrier, reduction of wrinkles, firming
R.F. 11.12.34	signs of aging skin, barrier disorders due to wrong cosmetic products, wrinkles	slight firming and improvement of skin condition	slight improvement, "might be due to cosmetic products": +	6 J* cm ² / 1 pulse	minor improvement of skin surface
B.S. 28.11.60	minor wrinkles, dry skin with microscopic barrier disorders	improved skin condition, firming	visible improvement and refreshed skin condition: +++	8 J* cm ² / 2 pulses	improved skin barrier

Tab. 1: Treatment results



Fig. 2a: Before the IPL/boswellia treatment



Fig. 2b: After the IPL/boswellia treatment




Fig. 3a: Barrier disorder before the treatment



Fig. 3b: Barrier disorder after the treatment

The condition of the skin before and after the treatment has been photo documented and a dermatoscopy (with 10 x magnification) was carried out. In addition, the visual appearance of the skin was evaluated with a magnifying glass according to current dermato-cosmetic criteria. Before and after the respective treatments the test persons were asked to subjectively assess their skin through a mirror.

Evaluation and discussion

The innovative antiaging treatment with IPL activated boswellia nanoparticles proved to be very effective from the subjective as well as the objective perspective. Looking in the mirror after the first session the test persons already rated the result of the treatment as “excellent”. The treatment was not only successful on the short term though, after 5 treatment sessions a long term improvement of the skin barrier, a reduction of the wrinkle depth and a firming of the skin could be observed (checkup after 4 weeks).

By using the energy of the IPL flash lamp the efficacy of boswellia nanoparticles against inflammatory skin diseases and barrier disorders with premature skin aging as already established in previous pilot studies could still be improved. Based on the chemical structure of the acetyl-keto-boswellic acid and the known infrared absorption spectrum it was extrapolated that triterpenes could be activated by high energy light. From a biological perspective so an improved efficacy of the treatment with boswellia extract should be observable.

The activation and improved efficacy could be verified and confirmed in the present study. Significant in this context were triterpenes which can be activated by absorption of light. Acetyl-keto-boswellic acid is a triterpene of the frankincense resin. However, the resin will not penetrate through the skin or in other words, once applied frankincense resin only sticks to the skin and the ingredients will not permeate into the deeper skin layers. Only by means of a specific manufacturing process called high

pressure homogenization which consists of encapsulating the substance into a shell of natural phospholipid, the so-called phosphatidylcholine, nanoparticles are produced which will fuse with the skin barrier and hence transport the boswellic acids into the deeper skin layers. There they can additionally be activated by IPL light energy in order to develop their specific effect as e.g. inhibiting the collagen degrading matrixmetalloproteinases and the inflammation mediators like 5-lipoxygenases.

Concluding, the study proved that the IPL treatment with light-activated boswellia nanoparticles is an innovative and very effective cosmetic therapy to rejuvenate the skin and treat inflammatory skin diseases.

References:

1. H.-U. Jabs, Behandlung aktinischer Keratosen mit einem neuen Weihrauchextrakt. Kosmetische Medizin 4 (2005) 4-5.
2. H.-U. Jabs, Aktinische Keratosen - eine Volkskrankheit. Ästhetische Dermatologie 4 (2005) 30-32.
3. H.-U. Jabs, Entzündliche Hauterkrankungen durch oxidativen und nitrosativen Stress. Ästhetische Dermatologie 3 (2008) 28-36.
4. H.-U. Jabs, Exotische Wirkstoffe - Schätze aus Fernost. Profi Kosmetik 07/08 (2008) 20-21.

Dr. Hans-Ulrich Jabs